OIPE
OCT 25 2004
PATENT & TRADEMARK OFFICE



FIGURE 1. Concentration of IFNγ, IL-4 and IL-5 in spleen cell supernatants of mice infected with *M. avium*, *S. mansoni* or both organisms. Splenocytes ($4x10^5$/well) were cultured *in vitro* for 48h at 37°C in 200 µl medium in the presence or absence of optimal concentrations of PPD or soluble schistosome egg antigen (SEA). Cytokine secretion was quantified by ELISA.

Figure 1

BEST AVAILABLE COPY



FIGURE 2. Concentration of IFNγ and IL-4 in granuloma cell supernatants of mice infected with *M. avium*, *S. mansoni* or both organisms. Granuloma cells ($4x10^5$/well) in 200 µl medium were cultured *in vitro* at 37°C for 48h in the presence or absence of the optimal concentration of PPD or soluble schistosome egg antigen (SEA). Cytokines secretion was quantified by ELISA.

Figure 2



FIGURE 3. Serum IgG1, IgE and IgG2a levels in mice infected with *M. avium*, *S. mansoni* or both (concurrent) as described above. Immunoglobulin concentration was determined by ELISA.

Figure 3



FIGURE 4. IL-4 can cure mice of a chronic *H. polygyrus* infection. Mice received three injections of an IL-4 complex starting 12 days before sacrifice. Data are means ±SE of multiple determinations.

Figure 4

MICE PREVIOUSLY COLONIZED WITH AN INTESTINAL HELMINTH DEVELOP ATTENUATED TNBS COLITIS



Fig 5

COLONIZATION WITH *H. POLYGYRUS* INHIBITS MUCOSA IFNγ REPONSIVENESS



Fig 6


H. POLYGYRUS BLOCKS
MUCOSA IL12 SYNTHESIS
250
200
150
100
50
0
IL12 (pg/ml)
Control
H. poly

Fig 7

COLONIZATION WITH *H. POLYGYRUS* PROMOTES MUCOSA IL10 PRODUCTION



Fig 8

BLOCKADE OF IL10R ENHANCES LPMC IFNγ PRODUCTION



Fig 9

COLONIZATION WITH *H. POLYGYRUS* PROMOTES MUCOSA PGE2 PRODUCTION



Fig 10

COLONIZATION WITH *H. POLYGYRUS* PROMOTES MUCOSA IL4, IL5 AND IL13 PRODUCTION



Fig 11

COLONIZATION WITH *H. POLYGYRUS* PROMOTES MUCOSA TGFβ PRODUCTION



Fig 12

T CELLS MAKE THE IFNγ IN THE INTESTINAL MUCOSA OF HEALTHY WT MICE



Fig 13

TRANSFER OF MLN CELLS FROM *H POLYGYRUS*-BEARING MICE INTO UNINFECTED WT MICE INHIBITS LPMC IFNγ RESPONSIVENESS



Fig 14

MLN T CELLS FROM MICE BEARING *H. POLYGYRUS* ENTER GUT MUCOSA WHEN TRANSFERRED INTO WT RECIPIENTS



Fig 15

PIROXICAM-INDUCED COLITIS IN IL10-/- MICE



Fig 16

H. POLYGYRUS REVERSES ESTABLISHED ACTIVE PIROXICAM-INDUCED IL10-/- COLITIS



Fig 17

H. POLYGYRUS BLOCKS LPMC IFNγ AND IL12 PRODUCTION IN IL10-/- COLITIS



Fig 18

H. POLYGYRUS AUGMENTS LPMC IL4 AND IL13 PRODUCTION IN IL10-/- COLITIS



Fig 19

MLN CELLS FROM IL10KO MICE COLONIZATION WITH *H. POLYGYRUS* INHIBIT ACTIVE IL10KO IBD



Fig 20

***H. POLYGYRUS* AUGMENTS MLN CELL EXPRESSION OF *Foxp3* mRNA AS MEASURED WITH REAL TIME RT-PCR**



Fig 21

***H. POLYGYRUS* REDUCES MLN CELL EXPRESSION OF *Smad7* mRNA AS MEASURED WITH REAL TIME RT-PCR**



Fig 22



Fig. : antigens - diagram of extraction procedures

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☑ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.
As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.